



# Histoacryl<sup>®</sup> LapFix

A REVOLUTION IN MESH FIXATION!

# Histoacryl® LapFix

## A REVOLUTION IN MESH FIXATION!

### ATRAUMATIC

Atraumatic mesh fixation with cyanoacrylate is a solution available for patients undergoing laparoscopic inguinal hernia surgery

- Non invasive method reported to be better tolerated by the patient<sup>1,2,3,4</sup>
- Low recurrence rate<sup>1,4,5,6,8,11,12,13,14</sup>
- Lower chronic pain<sup>6,8,12</sup> or no chronic pain compared to tackers<sup>7,11,13,15,16</sup>
- Less seroma compared to tackers<sup>9,10,11</sup>
- Faster return to work<sup>6</sup>

### SAFE

Atraumatic mesh fixation with cyanoacrylate is a solution proven to be as safe as the standard laparoscopic fixation with tackers

- Good biocompatibility and in vivo tolerance<sup>6,17,18,19</sup>
- No human or animal components<sup>21</sup>
- Strong fixation (prevents mesh dislocation)<sup>9,20</sup>
- Intrinsic bacteriostatic properties that help to reduce local septic complications<sup>5</sup>


### EASY

Atraumatic mesh fixation with cyanoacrylate is a very convenient solution that bring many advantages compared to tackers or sutures

- Fast and efficient polymerization<sup>6</sup>
- Storage at room temperature<sup>21</sup>
- Ready to use system, assembly in < 30 sec



In internal pre-clinical test<sup>25</sup>, a group of experienced surgeons n=6 evaluated the easiness of use



- Evaluate the easiness to assemble the cannula to the device (from 1 to 10).  
**Average score 9.33**
- Is the shape of the type adequate for the procedure (from 1 to 10)?  
**Average score 9.00**
- Will it possible to fix a bilateral hernia with this device? **100 % yes**
- Does the device look functional and useful for being used to fix meshes in laparoscopic Hernia repair procedures? **100 % yes**

## REFERENCES / STUDIES

- 1 Testini M, Lissidini G, Poli E, Gurrado A, Lardo A, Piccinni G. A single-surgeon randomized trial comparing sutures, N-butyl-2-cyanoacrylate and human fibrin glue for mesh fixation during primary inguinal hernia repair. *Can J Surg.* 2010;53(3):155-60.
- 2 Agresta F, Baldazzi GA, Ciardo LF, Trentin G, Giuseppe S, Ferrante F et al. Lightweight partially absorbable monofilament mesh (polypropylene/polyglactone 25) for TAPP inguinal hernia repair: initial experience. *Surg Laparosc Endosc Percutan Tech.* 2007;17(2):91-4.
- 3 Helbling C, Schlumpf R. Sutureless Lichtenstein: first results of a prospective randomised clinical trial. *Hernia.* 2003;7(2):80-4.
- 4 Nowobilski W, Dobosz M, Wojciechowicz T, Mionskowska L. Lichtenstein inguinal hernioplasty using butyl-2-cyanoacrylate versus sutures. Preliminary experience of a prospective randomized trial. *Eur Surg Res.* 2004;36(6):367-70.
- 5 Farouk R, Drew PJ, Qureshi A, Roberts AC, Duthie GS, Monson JR. Preliminary experience with butyl-2-cyanoacrylate adhesive in tension-free inguinal hernia repair. *Br J Surg.* 1996;83(8):1100.
- 6 Kukleta JF, Freytag C, Weber M. Efficiency and safety of mesh fixation in laparoscopic inguinal hernia repair using n-butyl cyanoacrylate: long-term biocompatibility in over 1,300 mesh fixations. *Hernia.* 2012;16(2):153-62.
- 7 Treepongkaruna SA, Subwongcharoen S. Novel technique of mesh fixation with cyanoacrylate in totally extraperitoneal laparoscopic repair: early experience. *J Med Assoc Thai.* 2012;95 Suppl 3:98-101.
- 8 Brügger L, Bloesch M, Ipaktchi R, Kurmann A, Candinas D, Beldi G. Objective hypoesthesia and pain after transabdominal preperitoneal hernioplasty: a prospective, randomized study comparing tissue adhesive versus spiral tacks. *Surg Endosc.* 2012;26(4):1079-85.
- 9 Lau, H. Fibrin sealant versus mechanical stapling for mesh fixation during endoscopic extraperitoneal inguinal hernioplasty: a randomized prospective trial. *Ann Surg.* 2005;242(5):670-5.
- 10 Topart P, Vandenbroucke F, Lozac'h P. Tisseel vs tack staples as mesh fixation in totally extraperitoneal laparoscopic repair of groin hernias: a retrospective analysis. *Surg Endosc.* 2005;19(5):724-7.
- 11 Jani K. Randomised controlled trial of n-butyl cyanoacrylate glue fixation versus suture fixation of mesh in laparoscopic totally extraperitoneal hernia repair. *J Minim Access Surg.* 2016;12(2):118-23.
- 12 Subwongcharoen S, Ruksakul K. A randomized controlled trial of staple fixation versus N-butyl-2-cyanoacrylate fixation in laparoscopic inguinal hernia repair. *J Med Assoc Thai.* 2013;96 Suppl 3:8-13.


STUDY	YEAR	PATIENT	PRODUCT	TECHNIQUE
Kukleta et al. <sup>6</sup>	2011	1336	BCA vs. tacks	TAPP
Mikhail et al. <sup>22</sup>	2012	198	BCA vs. suture	Plug in open technique
Brügger et al. <sup>8</sup>	2012	80	BCA vs. tackers	TAPP
Moreno Egea et al. <sup>15</sup>	2013	70	BCA vs. non absorbable suture	Lichtenstein/Rutkow and TEP
Subwongcharoen et al. <sup>12</sup>	2013	60	Histoacryl® vs. staples	TEP
Wang et al. <sup>13</sup>	2013	1027	No fixation vs. BCA vs. S vs. BCA + S	TAPP
Gutlic N et al. <sup>23</sup>	2016	1110	Permanent fixation (staplers/tackers) vs. BCA/no fixation	TEP
Filipovic-Cugura et al. <sup>14</sup>	2014	30	Histoacryl® vs. tackers	TEP
Golling et al. <sup>20</sup>	2014	28	Histoacryl® vs. tackers	TAPP and TEP
Burza et al. <sup>16</sup>	2014	70	BCA vs. tackers	TAPP
Garcia-Vallejo et al. <sup>24</sup>	2014	61	BCA	TEP
Jani K <sup>11</sup>	2016	251	BCA vs. absorbable suture	TEP

NR: not recorded, > better or faster, = equal, < smaller or lower, H: Histoacryl®, BCA: Butyl-cyanoacrylate, S: suture, T: tacks, F: fibrin glue, NS: no significance

- 13 Wang MG, Tian ML, Zhao XF, Nie YS, Chen J, Shen YM. Effectiveness and safety of n-butyl-2-cyanoacrylate medical adhesive for noninvasive patch fixation in laparoscopic inguinal hernia repair. *Surg Endosc.* 2013;27(10):3792-8.
- 14 Filipovic-Cugura J, Romic M, Misir Z, Filipovic N. Laparoscopic total extraperitoneal (TEP) inguinal hernia repair, comparison between postoperative pain using Hystoacril® Braun glue or Covidien ProTack™ fixation device for mesh fixation. *Hernia.* 2014;18(2):55-134.
- 15 Moreno-Egea A. Is sutureless hernia repair a safe option for treating abdominal Wall Hernias? A prospective study with a synthetic tissue adhesive (n-hexyl-alpha-cyanoacrylate). *Cir Esp.* 2013;91(4):243-9.
- 16 Burza A, Avantifiori R, Curinga R, Santini E, Delle Site P, Stipa F. Comparison between two different mesh fixation methods in laparoscopic inguinal hernia repair: tackler vs. Synthetic cyanoacrylate glue. *Minerva Chir.* 2014;69(6):321-9.
- 17 Eldabe Mikhail A, Palomo Luquero A, Reoyo Pascual JF, Seco Gil JL. Prosthetic material fixation in open inguinal hernioplasty: suture vs. synthetic glue. *Cir Esp.* 2012;90(7):446-52.
- 18 Shen YM, Sun WB, Chen J, Liu SJ, Wang MG. NBCA medical adhesive (n-butyl-2-cyanoacrylate) versus suture for patch fixation in Lichtenstein inguinal herniorrhaphy: a randomized controlled trial. *Surgery.* 2012;151(4):550-5.
- 19 Pascual G, Sotomayor S, Rodríguez M, Pérez-Köhler B, Kühnhardt A, Fernández-Gutiérrez M et al. Cytotoxicity of Cyanoacrylate-Based Tissue Adhesives and Short-Term Preclinical In Vivo Biocompatibility in Abdominal Hernia Repair. *PLoS One.* 2016;11(6):e0157920.
- 20 Golling M, Hofmann P, Hess C. Mesh fixation for TAP and TEP – First use of a new laparoscopic cyanoacrylate applicator. *Hernia* 2014;18(2):135-47.
- 21 Instruction for Use Histoacryl®.
- 22 Mikhail AE, Palomo AP, Reoyo JF, Seco JL. Prosthetic material fixation in open inguinal hernioplasty: suture vs. synthetic glue. *Ciresp* 2012;90(7):446-52.
- 23 Gutlic N, Rogmark P, Nordin P, Petersson U, Montgomery A. Impact of mesh fixation on chronic pain in total extraperitoneal inguinal hernia repair (TEP): a nationwide register-based study. *Ann Surg.* 2016;263(3):1199-206.
- 24 Garcia-Vallejo L, Couto-Gonzalez I, Concheiro-Coello P, Brea-Garcia B, Taboada-Suarez A. Cyanoacrylate surgical glue for mesh fixation in laparoscopic total extraperitoneal hernia repair. *Surg Laparosc Endosc Percutan Tech.* 2014;24(3):240-3.
- 25 Internal data. Survey on the handling of Histoacryl® LapFix, Barcelona, Spain, May 2016

EARLY PO PAIN	CHRONIC PAIN	RECURRENCE	OPERATION TIME	FOLLOW-UP
BCA<T due to different factors. Not used as evidence.	BCA<T	0.37 % vs. 2.3 %	NR	98 months
NS	2.9 % vs. 10.3 %	0 % vs. 1 %	NR	16 months
2 % vs. 14 %	4 % vs. 32 %	BCA = T	BCA = T	38 months
2.4 +/- 1 vs. 4.5 +/- 1.4 (VAS)	0 % vs. 20 %	0 % vs. 0 %	30 min vs. 70 min	15 months
1.6 +/- 1.33 vs. 2.35 +/- 1.32 (VAS)	16 % vs. 33 %	0 % vs. 3.3 %	NR	1 year
1.4 ± 0.6 vs. 1.3 ± 0.6 vs. 2.2 ± 0.9 vs. 2.2 ± 0.7 (VAS)	0 % vs. 0 % vs. 2.2 % vs. 2.1 %	0 % vs. 0 % vs. 0 % vs. 0 %	NR	19 months
0 % vs. 0.5 %, NS	8.7 % vs. 7.4 %, NS	1.5 % vs. 1.3 %, NS	40 min	33 months recurrent hernia 7.5 years
1.8 vs. 2.3	NR	0 % vs. 0 %	NR	1 month
NR	NR	NR	NR	NR
NR	0 % vs. 11.42 %	2.85 % vs. 2.85 %	BCA = T	24 months
59 % (24h), 34.4 %(48h), 6.6 %(>48h)	0 %	0 %	NR	29.7 months
BCA<S but not statistically significant	0 % vs. 0 %	0 % vs. 0 %	BCA>S	24 months

## ORDERING INFORMATION

DESCRIPTION		ARTICLE NO.	CONTENTS
<b>Histoacryl® Lap Mesh Fixation:</b> 1 unit with 2 Histoacryl® ampoules		1052008	2 x 0.5 ml 1 x LapFix applicator
<b>Histoacryl® Lap Mesh Fixation:</b> 5 units with 5 Histoacryl® ampoules		1050165	5 x 0.5 ml 5 x LapFix applicator
<b>Histoacryl® Blue</b>		1050044	5 units
<b>Optilene® Mesh</b> 7.5 x 15 cm (60 g/m <sup>2</sup> ; pore size 1.5 mm)		1065030	5 units
<b>Optilene® Mesh</b> 10 x 15 cm (60 g/m <sup>2</sup> ; pore size 1.5 mm)		1065040	5 units
<b>Optilene® Mesh</b> 15 x 15 cm (60 g/m <sup>2</sup> ; pore size 1.5 mm)		1065080	5 units
<b>Optilene® Mesh LP blue striped</b> 7.5 x 15 cm (36 g/m <sup>2</sup> ; pore size 1 mm)		1964715	5 units
<b>Optilene® Mesh LP blue striped</b> 10 x 15 cm (36 g/m <sup>2</sup> ; pore size 1 mm)		1964725	5 units
<b>Optilene® Mesh LP blue striped</b> 15 x 15 cm (36 g/m <sup>2</sup> ; pore size 1 mm)		1964705	5 units
<b>Optilene® Mesh Elastic blue striped</b> 10 x 15 cm (48 g/m <sup>2</sup> ; pore size 2.4 x 3.6 mm)		1964920	5 units



B. Braun Surgical, S.A. | Carretera de Terrassa, 121 | 08191 Rubí | Spain  
Phone +34 93 586 6200 | Fax +34 93 588 1096 | [www.bbraun.com](http://www.bbraun.com)

## AESCULAP<sup>®</sup> – a B. Braun brand

Aesculap AG | Am Aesculap-Platz | 78532 Tuttlingen | Germany  
Phone +49 7461 95-0 | Fax +49 7461 95-2600 | [www.aesculap.com](http://www.aesculap.com)

The main product trademark "Aesculap" and the product trademarks "Histoacryl" and "Optilene" are registered trademarks of B. Braun Melsungen AG and its subsidiaries.

Subject to technical changes. All rights reserved. This brochure may only be used for the exclusive purpose of obtaining information about our products. Reproduction in any form partial or otherwise is not permitted.